
# Geel 2000 Language Schools Math Department

Second term

Prep. 3



2022/2023

Name :	
Class:	

# Unit 1 Lesson 1 : Solving 2 equations

1)Complete:			
	of the two equat	ions: $x + y =$	0, $y - 5 = 0$ in $R \times R$
is			
2) The S.S (	of the two equati	ons: $x + 3y$	= 4, $3y + x = 1$ in R
xRis			
3) The S.S (	of the two equati	ons: 4 x + y =	6 , 8x + 2y = 12 in
R x R is			
4) If $L_1 \cap L_2$	$=\emptyset$ , then the S.S	of the two eq	<b>Juations which are</b>
represe	nted by the two S	St. lines $L_1$ and	d L <sub>2</sub> are
-	y = 5, x - y = 3		
-		represent the	e two equations x +
•	x + ay = 7		
_	allel, then a =		
-	-		o equations $x + 2y =$
	x + ky = 2, then		qual
<u>2)Choose t</u>	he correct answe	er:	
-	St. lines : $2 \times = 3$		
	dicular (b)c	oincident	(c)parallel
(d)intersec	eting		
a>-1			
2)The two s	traight lines : y	= x - 3 , $y =$	= x + 3 are
(a) parallel		erpendicular	(c)coincident
(d)intersec	eting		
0)/m/	C. II. O F	0 = 0	
	St. lines : 3 x + 5	y = 0, 5x - 3	y = 0 are
intersecting a			
(a) the original	in noint		(h) the first quadrant
(a) the orig	ın pomt		(b) the first quadrant
(c) the 2 <sup>nd</sup> qu	ıadrant		(d)the 4th quadrant
• •		s of the two e	quations: $x + y = 2$
and $y - 3 = 0$			1
(a) zero	(b) one	(c) two	(d) Three
	• •		

3) Find the solution set of the equations in R x R:

$$A) \qquad x + y = 5$$

$$, x-y=1$$

B) 
$$2x-y=3$$
,  $x+2y=4$ 

C) 
$$y = x + 4$$
 ,  $x + y = 4$ 

4) Find algebraically & graphically in R x R the S.S of each pair of the following equations:

1) 
$$x + 2y = 8$$
 ,  $3x + y = 9$ 

2) 
$$2 x + y = 1$$
 ,  $x + 2y = 5$ 

3) 
$$x + 2y = 8$$
 ,  $3x + y = 9$ 

5)What's the number of solutions of each pair of the following equations:

1) 
$$7x + 4y = 6$$
 ,  $5x - 2y = 14$  ( ......)

3) 
$$9 x + 6 y = 24$$
 ,  $3 x + 2 y = 8$  (.....)


6) Find the value of a and b:


1) a x + b y + 5 = 0 , 2 a x + b y - 1 = 0 given that (1,2) is a solution for two equations


2) If: 
$$f(x) = a x^2 + b$$
,  $f(1) = 5$ ,  $f(2) = 11$ 


Use your mind to choose the correct answer:

1)Which of the following graphs represents two equations of the first degree in two variables which have no common solution?









## Unit 1 Lesson 2 : Applications


1)The sum of two natural numbers is 42 and their
difference is 10
Find the two numbers ?
2)A rectangle is with a length more than its width by 4
cm . If the perimeter of the rectangle is 28 cm .Find the
area of the rectangle?
area of the rectangle.
3)Two supplementary angles , the twice of the measure
of their bigger equals seven times the measure of the
smaller . Find the measure of each angle ?
4)Two acute angles in a right – angled triangle , the
difference between their measures is 50. Find the
measure of each angle ?


5)A rectangle of perimeter 24 cm . If its length decreased by 4 cm and its width increased by 2 cm
became a square .Find the area of the square ?
6)If three times a number is added to twice a second
number the sum is 2, and if the first number is added to three times the second number the sum is 10
Find the two numbers ?
7)Two complementary angles, three times the measure
of one of them is more than half the measure of the
other by 18. Find the measure of each of them?

# Unit 1 Lesson 3: Solving 2<sup>nd</sup> degree

Find in R the S.S of each of the following equations using the general formula:  1) $x^2 + 7x + 2 = 0$ approximate the result to the nearest tenth
2) x <sup>2</sup> - 4 x + 1 = 0 approximate the result to the nearest 2 decimals
3) Graph the function $f:f(x)=x^2-2x$ in the interval $[-1,3]$ , from the graph find the S.S of the equation : $x$ - $2x = 0$ ?
4) Graph the function $f: f(x) = x^2 - 4x + 3$ on the interval [-1,5] and from the graph, find: a) The minimum value of the function b) The equation of the axis of symmetry c) The S.S of the equation $f(x) = 0$

2) From following figures the S.S of the equation f(x) = 0 in R is





-3 - Choose the correct answer from the given ones

(1) the curve of the function f such that  $f(x) = x^2 - 3x + 2$  cuts xaxis at the two points......

$$(c) (-2,0), (-1,0)$$

$$(d)(2,0),(-1,0)$$

(2) The solution set of the equation  $2x^2 + 5x = 0$  ......

(a) 
$$\{0,5\}$$
 (b)  $\{0,\frac{-5}{2}\}$ 

$$(c) \{ 2, 5 \}$$

(d) 
$$\varphi$$

(3) The solution set of the equation  $x^2 - 4x + 4 = 0$  .....

(a) 
$$\{-2,2\}$$
 (b)  $\{4,1\}$ 

(4) The solution set of the equation  $x^2 + 5 = 0$  ......

(a) 
$$\{\sqrt{5}, -\sqrt{5}\}$$
 (b)  $\{-\sqrt{5}\}$  (c)  $\{\sqrt{5}\}$ 

(b) 
$$\{-\sqrt{5}\}$$

(c) 
$$\{\sqrt{5}\}$$

- (5) In the equation:  $ax^2 + bx + c = 0$ , if  $b^2 4a c > 0$ , then the number of roots equals:....
- (a) 1

(b) 2

(c) 0

(d) underdetermined

4- Find the solution set for each pair of the following equations by using the formula: (1)  $x^2 - 2x - 4 = 0$  knowing that  $\sqrt{5} \approx 2.24$ (2)  $x^2 = 2 (x + 6)$  knowing that  $\sqrt{52} \approx 7.2$ (3)  $(x-1)^2 = 10$  knowing that  $\sqrt{10} \approx 3.16$ (4)  $x^2$  - 2 (x+3) = 0knowing that  $\sqrt{7} \approx 2.65$ (5)  $(x-3)^2 - 3(x-3) + 1 = 0$  knowing that  $\sqrt{5} \approx 2.24$ (6)  $1 - \frac{2}{x} = \frac{2}{x^2}$  (where  $x \neq 0$ ) knowing that  $\sqrt{3} \approx 1.73$ 

 $(7) 9x^2 - 24x + 16 = 0$ 

(8)  $x^2 = 2 (x - 6)$ (9)  $x + \frac{4}{x} + 1 = 0$  (where  $x \neq 0$ ) (10) if  $x^4 + 2x^2 - 1 = 0$ 

#### Unit 1 Lesson 4: 1<sup>st</sup> & 2<sup>nd</sup> degree

#### 1)Complete:

1) If 
$$x^2 - y^2 = 40$$
,  $x + y = 10$ , then  $x - y = \dots$ 

2) The S.S of the two equations: 
$$x - y = 0$$
,  $xy = 9$  in RxR is ...........

3)The ordered pair which satisfies each of the two equations

$$xy = 2$$
,  $x - y = 1$  is ......

# 2- Find in R x R the S. S of each pair of the following equations:

1) $x - y = 0$	$2 x^2 - y^2 = 9$	
----------------	-------------------	--



2) 
$$x - y = 1$$
,  $x^2 + y^2 = 25$ 

3) 
$$y - x = 3$$
 ,  $x^2 + y^2 - xy = 13$ 

......

cm <sup>2</sup> . Find its two  Dimensions
4)For a rhombus, the difference between the lengths of its diagonals equals 4 cm and its perimeter is 40 cm. Find the lengths of the diagonals.
5)The sum of two real numbers is 9 and the difference between their squares equals 45 .Find the two numbers .
6)A length of a rectangle is 3 cm more than its width and its area is 28 cm <sup>2</sup> . Find its perimeter
7)A right angled – triangle of hypotenuse length 13 cm . and its perimeter is 30 cm . Find the lengths of the other two sides .

8) The sum of two real positive numbers is 17 and their
product is 72

#### Find the two numbers

.....

#### 9) Complete:

- 1) If the curve of the quadratic function does not intersect the x-axis at any point, then the number of solutions of the equation f(x) = 0 in R is ......
- 2) If the curve of the quadratic function intersects the x-axis at the two points (2,0), (-3,0), then the solution set of the equation f(x) = 0 in R is ......
- 3) If  $x \in \mathbb{R}$ , then the equation  $x^2 + x + 1 = 0$  has ..... roots.
- 4) The curve of the function  $f: f(x) = x^2 5x$  intersects the x- axis at the two points ....
- 5) If the S.S of the equation :  $4 x^2 + 4 x + k = 0$  is  $\{\frac{-1}{2}\}$ ,  $k = \dots$
- 6) If : x = 3 is one of the solutions for the equation :  $x^2 + b x 12 = 0$ , the  $b = \dots$

### **Model Exam on Unit One**

(1) The degr	ee of the equa	tion 3x + 4y + xy	= 5 is
(a) zero	(b) first	(c) second	(d) third
(2) one soluti	on of the equa	tion $x^2 - y^2 = 3$ in	R may be
(a) (1,-2)	<b>(b)</b> ( <b>-2</b> , 1 )	(c) (1,2)	(d)(-1,-2)
(3) the ordere xy = 2 & x -y	_	isfies both of the	two equations
(a) (1,2)	(b) (2,1)	(c) (1,1)	(d)(2,-1)
(4) The soluti	on set of the tv	wo equations: x =	= y, x y = 1
is (a) {(1,1)} (c) {(1,-1)}		(b) {( -1 , -1 (d){( -1 , -1	(1)} (1,1)} (1,1)}
(5) The solution	on set of the tw	vo equations: x -	y = 0, x y = 9
is (a) {(0,0)} (c) {(3,3)}			(3,3)}
(6) One soluti may be		tions: $x - y = 2$ ,	x +y = 20 in K
(a) (-4,2)	(b) (2,-4)	(c)(3,1)	(d)(4,2)

2 - Find the solution set for each pair of the following equations

$$(1) x + 1 = 0$$

$$x^2 + y^2 = 17$$

(2) 
$$x - 2 = 0$$

$$x^2 + xy + y^2 = 7$$

$$(3) x - y = 0$$

$$, xy = 1$$

$$(4) x + y = 0$$

$$2x^2 - y^2 = 4$$

$$(5) x - 2y = 0$$

$$x^2 - y^2 = 3$$

$$(6) x - y = 1$$

$$x^2 + y^2 = 25$$

$$(7) y = x - 5$$

$$x^2 - 2xy = 16$$

(8) 
$$y - x = 3$$

$$x^2 + xy - 4 = 0$$

$$(9) x - 2y - 1 = 0$$

$$x^2 - xy = 0$$

$$(10) y + 2x = 7$$

$$2x^2 + x + 3y = 19$$

3 - Applications

(1) If the sum of two integer numbers is 3, and the sum of their square is 5, find the two numbers.

(2) Two numbers one of them is the additive inverse of the other, and the sum of their squares is 2, find the two numbers.

(3) If the difference between two numbers is 5, and their product is 36, then find the two numbers.

(4) ) If the sum of two integer numbers is 9, and the difference between their square is 27 find the two numbers.

14

- (5) Find the number which is formed from two digits, if the units digit is twice the tens digit, and if the product of the two digits equals half the original number.
- (6) The length of a rectangle is 3 more than its width, and its area is  $28\text{cm}^2$ . find its perimeter.
- (7) Find the two dimensions of a rectangle if its perimeter is 24 cm. and its area is 35 cm<sup>2</sup>.
- (8) Find the two dimensions of a rectangle if its diagonal of length 5 cm. and its perimeter is 14 cm.
- (9) The hypotenuse of a right angle triangle is 13 cm. and its perimeter is 30cm. find the length of the other two sides,
- (10) The difference between the lengths of the two rhombus's diagonals is 4cm. and its perimeter is 40 cm. Find the length of each diagonal.
- 4- Answer the following questions:
- (1) Graph the function f where  $f(x) = x^2 3x + 2$ ,  $x \in [-1, 4]$  then from the

Graph find.

- (a) The vertex point of the curve
- (b) The maximum or minimum value of the function f.
- (c) The solution of the equation  $x^2 3x + 2 = 0$
- (2) Graph the function f where  $f(x) = x^2 4x 2$ ,  $x \in [-1, 5]$  then from the

Graph find the solution set of the equation f(x) = 0

#### Unit 2 Lesson 1 : Set of zeroes

1)Determine the set of zeroes of the polynomial function which are defined by the following rules in R:

1) 
$$f(x) = 4x + 12$$

$$2) f(x) = x^2 - 5x$$

$$3) f(x) = 6 x^3 - 18 x^2$$

4) 
$$f(x) = x^2 + 25$$

$$5) f(x) = (x-2)(x+3) + 4$$

2)Complete:

- 1) The set of zeroes of the function f:f(x) = -4x is
- 2) The set of zeroes of the function  $f : f(x) = 8x^2$  is ......
- 4) The set of zeroes of the function f: f(x) = 0 is ......

- 5) The set of zeroes of the function  $f: f(x) = x^2 49$  is
- 6) The set of zeroes of the function  $f: f(x) = x^2 + kx + 1$  is  $\emptyset$ , then k may equal ......

7) If 
$$z(f) = \{5\}$$
,  $f(x) = x^3 - 3x^2 + a$ , then  $a = \dots$ 

8) If 
$$z(f) = \{1, -2\}$$
,  $f(x) = x^2 + x + a$ , then  $a = \dots$ 

9) If the set of zeroes of the function f where  $f(x) = x^2 + k$  is  $\{6, -6\}$  Then  $k = \dots$ 

#### Complete the following:

- (1) The set of zeroes of f where f(x) = x 5 is ......
- (2) The set of zeroes of f where f(x) = 4 is ......
- (3) The set of zeroes of f where  $f(x) = x^2 + 9$  is ......
- (4) The set of zeroes of f where  $f(x) = 4x^2 9$  is ......
- (5) The set of zeroes of f where  $f(x) = (x-5)^2$  is ......

#### Unit 2 Lesson 2: Equality of functions equations

#### 1)Determine the domain of each the algebraic fractional functions:

1) 
$$n(x) = \frac{x-5}{6 \ x}$$

2) n(x) = 
$$\frac{x^2 + 2}{x^2 - 81}$$

3) n(x) = 
$$\frac{x^2 - 1}{x^2 + 1}$$

4) n(x) = 
$$\frac{6x}{6x^3 - 54x}$$

#### 2) Find the common domain of the following algebraic fractions:

1) 
$$\frac{7x}{x^3 - 27}$$
,  $\frac{x + 12}{x^2 - 144}$ 

$$(2)^{\frac{x^2+x+1}{2x^4}}, \frac{x^2-1}{x^2-x}$$

3) 
$$\frac{x+5}{2}$$
,  $\frac{3}{x^2-25}$ ,  $\frac{3x}{x^2-5x}$ 

#### 3)Complete:

- 1) The domain of the algebraic fraction function is R the set of .....
- 2) The domain of the function  $f:f(x) = \frac{1}{x^3}$  is ......
- 3) The domain of the function  $f:f(x) = \frac{x^3 4x}{2x + 4}$  is......
- 4) If  $n(x) = \frac{x}{x^2 + 4}$ , then  $n(x) \in R$  for each  $x \in \dots$
- 5) If n (x) =  $\frac{7}{x+a}$  and the domain of the function n is  $R - \{-2\}$ , a = ....
- 6) If the function  $f:f(x) = \frac{x-5}{x^2-a}$  has the domain  $R-\{-5, 5\}$ , a = .....

4) Reduce each of the following algebraic fractions to the simplest form showing the domain of each of them:

1) n(x)=
$$\frac{x^2-4x}{x^2-16}$$

2) 
$$n(x) = \frac{x^2 - 6x + 9}{2x^3 - 18x}$$

5) In each of the following, Show that 
$$n_1 = n_2$$
 or not:  
1)  $n_1(x) = \frac{x^3 - 1}{x^3 + x^2 + x}$ ,  $n_2(x) = \frac{(x - 1)(x^2 + 1)}{x^3 + x}$ 

2) $n_1(x) = \frac{2 x}{2 x + 8}$ ,	$n_2(x) = \frac{x^2 + 4x}{x^2 + 8x + 16}$
6)In each of the followin	g, Show that $n_1 = n_2$ or not
1)n <sub>1</sub> (x) = $\frac{x^2 - 9}{x^2 + 4x + 3}$	, $n_2(x) = \frac{x-3}{x+1}$
2) $n_1(x) = \frac{2}{2x - 10}$	$n_2(x) = \frac{5+x}{x^2-25}$
3)n <sub>1</sub> (x) = $\frac{x^2 - 3x}{x^2 - 6x + 9}$	$n_2(x) = \frac{x^2 + 3 x + 9}{x^3 - 27}$
$3)n_1(x) = \frac{x^2 - 3x}{x^2 - 6x + 9}$	
7)Reduce each of the follow simplest form showing the	, $n_2(x) = \frac{x^2 + 3 x + 9}{x^3 - 27}$
7)Reduce each of the follow	, $n_2(x) = \frac{x^2 + 3 x + 9}{x^3 - 27}$
7)Reduce each of the follow simplest form showing the	, $n_2(x) = \frac{x^2 + 3 x + 9}{x^3 - 27}$

#### Unit 2 Lesson 3: Operations on functions

# 1)In each of the following find n(x) in the simplest form showing the domain of n:

1) n (x) = 
$$\frac{x}{x^2 + 2x} + \frac{x + 4}{x + 2}$$

2)n(x)=
$$\frac{x}{x-4} - \frac{x+4}{x^2-16}$$

3) n (x) = 
$$\frac{x^2 + x}{x^2 - 1}$$
 -  $\frac{5 - x}{x^2 - 6 x + 5}$ 

4) n(x) =  $\frac{x-6}{2 x^2 - 15 x + 18} - \frac{x-5}{15-13 x + 2 x^2}$ 

5) n(x) = 
$$\frac{3 x + 15}{x^2 + 7 x + 10} + \frac{2 x^2 - 3 x - 2}{x^2 - 4}$$

6) n(x) = 
$$\frac{x^2 + 3 x + 2}{x^2 - 9}$$
 x  $\frac{x - 3}{x + 2}$ 

7) n(x) = 
$$\frac{x+2}{x^2-4}$$
 x  $\frac{2x-4}{x-3}$ 

8) n(x) = 
$$\frac{x^2 + 2x - 3}{x + 3} \div \frac{x^2 - 1}{x + 1}$$

9) n (x) = 
$$\frac{x^3 - 8}{x^2 + x - 6}$$
 ÷  $\frac{x^2 + 2x + 4}{2x + 6}$ 

.....

10) n(x) = 
$$\frac{x^2 - 2x - 15}{x^2 - 9}$$
 ÷  $\frac{2x - 10}{x^2 - 6x + 9}$ 

11) n (x) = 
$$\frac{x^2 + 2x}{x^3 - 27} \times \frac{x^2 + 3x + 9}{x + 2}$$
 then find n(6), n(-2)if it possible

2)Complete:

- 1) If  $\frac{a}{b}$ ,  $\frac{c}{d}$  are two algebraic fractions, then  $\frac{a}{b} + \frac{c}{d} =$
- 2) The domain of n : n (x) =  $\frac{3 x + 4}{x^2 + 25} + \frac{x 2}{x^2 + 7}$  is
- 3)If :  $x \in R \{2\}$ , then:  $\frac{x}{x-2} + \frac{2}{2-x} = \dots$
- 4) The additive inverse of the fraction:  $\frac{x+7}{x-5}$  is .....
- 5) The function f where  $f(x) = \frac{x-2}{x-5}$  has additive inverse if the domain is ......
- 6) If  $n(x) = \frac{x^2 5x + 6}{5x}$ , then the domain of  $n^{-1}$  is

- 9) If the function  $n : n(x) = \frac{2 x}{x 2}$  has a multiplicative inverse, then the domain of n is ......

10) If the algebraic fraction  $\frac{x-a}{x-3}$  has a multiplicative inverse which is  $\frac{x-3}{x+2}$ , then a= ......

# 3)In each of the following find n(x) in the simplest form showing the domain of n:

1) n (x) = 
$$\frac{2x+6}{x^2+x-6} - \frac{x^2-6x}{x^2-8x+12}$$

2) n(x) = 
$$\frac{3x^2 + 6x}{x^2 - 4} + \frac{6}{2 - x}$$

3) n(x) = 
$$\frac{3 x - 15}{x + 3}$$
 x  $\frac{4 x + 12}{5 x - 25}$ 

4)n(x) = 
$$\frac{x^2 - 2 x - 3}{5 x - 135}$$
x  $\frac{5 x^2 + 15 x + 45}{x + 1}$ 

#### 4) find the answer of the following:

1)If: n(x) = 
$$\frac{x^2 - 2x}{(x - 2)(x^2 + 2)}$$

First: find  $n^{-1}(x)$  and identify its domain

Second: if  $n^{-1}(x) = 3$  what is the value of x?

2)If 
$$f(x) = \frac{x^2 - 49}{x^3 - 8} \div \frac{x + 7}{x - 2}$$
, then find  $n(x)$  in the simplest form and Identify its domain and find  $f(1)$ 

identity its domain and inid f(1)

3)If n (x) = 
$$\frac{x^3 + 3x^2 + 2x}{x^2 + 2x}$$
 find n<sup>-1</sup> (x) in the simplest form showing the domain of n<sup>-1</sup>, then find n<sup>-1</sup> (-2) if it is possible

4) Put the function 
$$f: f(x) = \frac{3 x + 6}{4 x - x^2 + 5} \times \frac{x - x^2 + 20}{7 x + 14}$$
 in the simplest form and identify its domain , then find if it is possible the value of  $f(-2)$  ,  $f(2)$ 

5)If 
$$f(x) = \frac{x^2 - 2x - 15}{x^2 - 9} \div \frac{x^2 - 25}{x^2 - 3x}$$
 find  $f(x)$  in the

simplest form showing the domain of  $f(a) = \frac{1}{3}$  find the value of a

#### **Model Exam on Unit two**

1) Choose the correct answer from the given ones

(1) The function f where  $f(x) = \frac{x-2}{x^3+27}$ , then the domain of its multiplicative inverse is R - .....

(a) {2}

(b)  $\{-3, 2\}$  (c)  $\{2, -3, 3\}$  (d)  $\{3, -3\}$ 

(2) If the function f where  $f(x) = \frac{x^2 - 9}{x}$ , has a multiplicative inverse, then their common domain is .....

(a)  $R - \{0\}$ 

(b)  $R - \{0, 3\}$ 

(c)  $R - \{0, 3, -3\}$  (d) R

(3) If  $N(x) = \frac{x-1}{x-2}$ , the domain of  $N^{-1}(x)$ ......

(b)  $R - \{1\}$  (c)  $R - \{2\}$  (d)  $R - \{1, 2\}$ 

(4) the function f where  $f(x) = \frac{x-2}{x-5}$ , has a multiplicative inverse if its domain is .....

(a) **R** 

(b)  $R - \{5\}$  (c)  $R - \{2\}$  (d)  $R - \{2, 5\}$ 

(5) the function f where  $f(x) = \frac{x-2}{x-5}$ , has a multiplicative inverse if the domain is .......

(a)  $R - \{2\}$  (b)  $R - \{5\}$  (c)  $R - \{-2, 2\}$  (d)  $R - \{0, 1\}$ 

(6) If  $N(x) = \frac{1}{x} - \frac{3}{x}$ , then  $N^{-1}(x) = \dots$ 

(a)  $x - \frac{x}{3}$  (b)  $\frac{2}{x}$ 

 $(c)\frac{-x}{2}$ 

 $(d)\frac{x}{2}$ 

(7) The domain of the function f where  $f(x) = \frac{x(x+2)}{x^2-4}$  is ......

(a) **R** 

(b)  $R - \{-2, 2\}$  (c)  $R - \{2, 0\}$  (d)  $R - \{2\}$ 

- (8) The domain of the function f where  $f(x) = \frac{x-3}{2}$  is ......
  - (b)  $R \{0\}(c) R \{-1, 0\} (d) R \{0, 1\}$ (a) **R**
- (9) The domain of the function f where  $f(x) = \frac{x-7}{3(x+1)}$  is ......
  - (b)  $R \{1\}$  (c)  $R \{-1, 3\}$  (d)  $R \{-1\}$ (a) **R**
- (10) The domain of the function n where  $n(x) = \frac{x-1}{x+2} + \frac{x-2}{x+1}$  is.....
- (a)  $R \{ -1 \}$

(c)  $R - \{-1, -2\}$ 

- (b) R { -2} (d) R { -1, -2, 1, 2}
- (11) If  $n(x) = \frac{2x}{x^2 x + 2}$ , then n(-1)......
- $(a)^{\frac{-1}{2}}$
- (b)  $\frac{1}{2}$  (c)  $\frac{2}{3}$
- $(\mathbf{d}) 2$
- (12) The function f where  $f(x) = \frac{x+2}{x-2}$ , then the domain of its multiplicative inverse is ....
  - (a) **R**

- (b)  $R \{2\}$  (c)  $R \{-2\}$  (d)  $R \{-2, 2\}$
- (13) The domain of the function n where  $n(x) = \frac{x-2}{x+3} \frac{3x}{x-1}$  is.....
  - (a)  $R \{0, 2\}$  (b)  $R \{-3, 1\}$  (c)  $R \{2, 3\}$  (d)  $R \{-3, 2\}$
- (14) The simplest form of the function  $n(x) \frac{x}{x-3} \div \frac{3x}{x^2-9}$  is ...
- $(a)\frac{x}{x-2}$

- (b)  $\frac{x}{x+3}$  (c)  $\frac{x+3}{2}$  (d)  $\frac{x-3}{2}$
- (15) The domain of the multiplicative inverse of the fraction  $\frac{x+7}{x-2}$  is ....
- (a) **R**

- (b)  $R \{2\}$  (c)  $R \{-7\}$  (d)  $R \{-7, 2\}$

(16) The additive inverse of the fraction  $\frac{3}{x^2+1}$  is..... (a)  $\frac{-3}{x^2+1}$  (b)  $\frac{x^2+1}{3}$  $(c)^{\frac{x^2+1}{2}}$   $(d)^{\frac{3}{x^2-1}}$ (17) If  $f(x) = \frac{x^2-9}{x+h}$ , f(4) = 1, then  $b = \dots$ (c) 3 (d) -3(a) - 7(b) 7 (18) If  $N(x) = \frac{x-2}{x^2-x-6}$ , then the domain of  $N^{-1}(x)$  R – ...... (c)  $\{-2,2\}$  (d)  $\{-2,2,3\}$ (a)  $\{2\}$  (b)  $\{-2,3\}$ (19) The simplest form of the function n(x)  $\frac{x+1}{x-1} + \frac{1-x}{x-1}$   $x \ne 1$ is .....  $(d) \frac{2}{(x-1)^2}$  $(b)\,\frac{2}{2x-2}$ (a) zero (20) The set of zeroes of f where  $f(x) = (x-1)^2(x+2)$  is... (b)  $\{1, -2\}$ (c) { -1, 2} (d) { -1, -2} (a) { 1, 2 } 2) Answer the following questions (1) Simplify each of the two algebraic  $n_1(x) = \frac{x^2 - 1}{x^2 - x}, n_2(x) = \frac{2x - 6}{x^2 - 5x + 6}$ (2) Simplify the function where  $n(x) = \frac{3x}{x^2 - 2x} - \frac{12}{x^2 - 4}$ , showing its domain.

(3) Simplify the function where $n(x) = \frac{x^2-1}{x^2+3x+2} - \frac{x^2-x}{x^2+2x}$ , showing its domain.
(4) Find n in its simplest form where $n(x) = \frac{x}{4} + \frac{-2}{x+2}$ , showing its domain.
(5) If the domain of the fraction n where $n(x) = \frac{b}{x} + \frac{9}{x+a}$ is R –{ 0 , 4 } ,n(5)= 2 Find the value of a and b
(6) Find n in its simplest form where $n(x) \frac{3x^2+6x}{x^2-4} \times \frac{x-2}{2x+6}$ , showing its domain.
(7) Find n in its simplest form where $n(x) \frac{x+3}{(x-2)(x+7)} \div \frac{2x+3x}{2x+14}$ , showing its domain.

(8) If $n_1(x) = \frac{x^2}{x^3 - x^2}$ , $n_2(x) = \frac{x^1 + x^2 + x}{x^4 - x}$ , prove that $n_1 = n_2$ .
(9) Find n in its simplest form , showing its domain . where
1) $\mathbf{n}(\mathbf{x}) = \frac{\mathbf{x}}{\mathbf{x}+1} + \frac{2\mathbf{x}^2}{\mathbf{x}^3 - \mathbf{x}}$
2) $\mathbf{n}(\mathbf{x}) = \frac{\mathbf{x}-1}{\mathbf{x}^2-1} \div \frac{\mathbf{x}^2-5\mathbf{x}}{\mathbf{x}^2-4\mathbf{x}-5}$
(10) Find n in its simplest form where
(10) Find n in its simplest form where $f(x) = \frac{3x^2 - 6x}{x^2 - 4} \times \frac{x^2 + 3x + 2}{x^2 + x}$
$f(x) = \frac{3x^2 - 6x}{x^2 - 4} \times \frac{x^2 + 3x + 2}{x^2 + x}$
$f(x) = \frac{3x^2 - 6x}{x^2 - 4} \times \frac{x^2 + 3x + 2}{x^2 + x}$
$f(x) = \frac{3x^2 - 6x}{x^2 - 4} \times \frac{x^2 + 3x + 2}{x^2 + x}$
$f(x) = \frac{3x^2 - 6x}{x^2 - 4} \times \frac{x^2 + 3x + 2}{x^2 + x}$ (11) Find the common domain of $f_1$ , $f_2$ to be equal such that: $f_1(x) = \frac{x^2 + x - 12}{x^2 + 5x + 4} \qquad , \qquad f_2(x) = \frac{x^2 - 2x - 3}{x^2 + 2x + 1}$
$f(x) = \frac{3x^2 - 6x}{x^2 - 4} \times \frac{x^2 + 3x + 2}{x^2 + x}$ $(11) \text{ Find the common domain of } f_1, f_2 \text{ to be equal such that:}$ $f_1(x) = \frac{x^2 + x - 12}{x^2 + 5x + 4} \qquad , \qquad f_2(x) = \frac{x^2 - 2x - 3}{x^2 + 2x + 1}$

(12) Find n in its simplest form, showing its domain. where 1)  $f(x) = \frac{x^2 - x}{x^2 - 1} + \frac{x - 5}{x^2 - 6x + 5}$ 2)  $f(x) = \frac{x^3 - 1}{x^2 - 2x - 1} \times \frac{2x - 2}{x^2 + x + 1}$ (13) If  $f(x) = \frac{x^2 - 49}{x^3 - 8} \div \frac{x + 7}{x - 2}$  find f in its simplest form, showing its domain. Then calculate f(1). (14) Find the common domain of  $f_1$ ,  $f_2$  to be equal such that:  $f_1(x) = \frac{x^2 + 3x + 2}{x^2 - 4}$ ,  $f_2(x) = \frac{x^2 - 1}{x^2 - 3x + 2}$ (15) Find n in its simplest form, showing its domain. where  $f(x) = \frac{3x}{x^2 - x - 2} + \frac{x - 1}{1 - x^2}$ 

#### **Unit 3:** Operations on events

#### 1- Complete the following

(1) The two events are said to be mutually exclusive if

$$A \cap B = \dots$$

- (2) If the probability that the event A occurs is 75%, the probability of non occurrence of this event is ......
- (3) If A is an event, P(A) = 0 then A is......
- (4) If A' is the complement event of A then  $A \cup A' = ....$ ,

$$A \cap A' = \dots$$

- (5) The probability of the sure event equals ......
- (6) The probability of the impossible event equals ......
- (7) When a regular die is tossed once, then the probability of getting an even number is ......
- (8) When a regular die is tossed once, then the probability of getting a head is....
- (9) If A, B are two mutually exclusive events, P(A) = 0.2,

$$P(B) = 0.3 \text{ then } P(A \cup B) = \dots$$

- (10) If A ,B are two mutually exclusive events of a random experiment , $P(A \cap B) = \dots$
- (11) If  $A \subset S$  of a random experiment, P(A) = P(A')

Then 
$$P(A) = ....$$

 $(12)\ (10)\ If\ A\ ,B$  are two mutually exclusive events of a random experiment,

$$P(A) = \frac{1}{4}$$
,  $P(A \cup B) = \frac{5}{12}$  then  $P(B) = .....$ 

#### 2]Find the result in each:

1]If A and B are two events in the sample space for a random experiment Where P ( A ) = 0.7 , P ( B ) = 0.6 and P ( A  $\cup$  B ) = 0.9 , **then find :** 

2]If A and B are two events in the sample space of a random experiment and there is : P(A) =  $\frac{1}{3}$ , P(B) =  $\frac{1}{2}$ ,

$$P(A \cap B) = \frac{1}{5}Find : P(A \cup B)$$

3]A ball is drawn randomly from 25 identical balls of the same volume and weight, 10 of them are red, 8 are white and the rest are green .**Find the Probability that the**drawn ball is:1) white

2)green or white

4]If A, B are two events from a sample space of a random experiment,

And 
$$P(A) = 0.8$$
,  $P(B) = 0.3$ ,  $P(A \cup B) = 0.9$   
**Find:** 1)  $P(A \cap B)$ .....

numbered from 1 to 30, find the probability that the number on the drawn card is:
First:Divisible by 4 second: A prime number
6]A ball is drawn randomly from 25 identical balls of the same volume and weight, 10 of them are red, 8 are white and the rest are green, find the Probability that the drawn ball is:  First:Whitesecond: Green or white
7] If A , B are two events in a random experiment , $P(A) = 0.7$ , $P(B) = 0.6$ and $P(A \cap B) = 0.4$ Find the <b>probability of :</b> 1) Non occurrence of the event A 2) Occurrence of one event without the other .
8]A classroom consists of 40 students, 30 of them play football, 20 play Basketball and 15 play football and basketball, if a student is chosen randomly. <b>Find: First:</b> The probability that this student is playing one of the two games at least
<b>Second :</b> The probability that this student is playing only one of the two games .
9]If $P(A) = 4 P(A)$ , then $P(A) =$

#### **Models Exams in Algebra and statistics**

#### Model (1)

#### First: Complete the following

- (1) If x is a negative number, the greatest number of the following: 5 + x, 5 x, 5x, and  $\frac{5}{x}$  is .........
- (2) If  $x \in R\text{-}\{0,\!1\}$  , then  $\frac{1-x}{x} \div \frac{x-1}{x}$  in the simplest form equals......
- (3) If A ,B are two events in a random experiment, and  $B \subset A$ , then  $P(A \cap B) = ....$
- (4) If the sum of two positive numbers is 4, and the sum of their squares is 10, Then the two numbers are .......
- (5) If the solution set of the equation  $x^2 + mx + 9 = 0$  is  $\{-3\}$ , then  $m = \dots$
- (6) If  $n(x) = \frac{x+7}{x-2}$ , then the domain of  $n^{-1}$  is ......

Second: Choose the correct answer from the given ones

- (1) the common domain of the two fractions  $\frac{2}{x-3}$ ,  $\frac{7}{x-6}$  is
- (a) **R**

- (b)  $R \{3\}$  (c)  $R \{6\}$  (d)  $R \{3, 6\}$
- (2) The probability of the impossible event equals ......
  - (a)  $\varphi$
- (b) zero
- (c) ]
- (d) -1

- (3) If 2x = 1, then  $\frac{1}{5}x = \dots$ 
  - (a)  $\frac{2}{5}$  (b)  $\frac{1}{5}$

- (4) If  $x^2 y^2 = 2(x + y)$  such that  $x + y \neq 0$  then
- $\mathbf{x} \mathbf{y} = \dots$

- (c) 6
- (d) 8
- (5) The set of zeroes of the function f where  $f(x) = \frac{x-3}{x+2}$  is
  - (a) {zero}
- (b) { 3 }
- (c)  $\{-2\}$  (d)  $\{3, -2\}$
- (6) The solution set of the two equation: -2y = 1, 3x + y = 10is .....
- (a)  $\{(5,2)\}$  (b)  $\{(3,1)\}$  (c)  $\{(4,2)\}$  (d) $\{(3,1)\}$

[3](a) solve the equation  $3x^2 = 5x + 4$  approximate to the nearest two decimals

(b) Find n in the simplest form showing its domain where

$$\mathbf{n}(\mathbf{x}) = \frac{\mathbf{x}^2 + 2\mathbf{x} + 4}{\mathbf{x}^3 - 8} + \frac{\mathbf{x}^2 + \mathbf{x} - 2}{\mathbf{x}^2 - 4}$$

- [4] (a) Graph the function f where  $f(x) = x^2 2x + 1$  over the interval [-2, 4] then from the graph find the solution of the equation  $x^2 2x + 1 = 0$
- (b) if  $n(x) = \frac{x^2 + 3x}{x^2 9} + \frac{2x}{x + 3}$  find n in its simplest form showing its domain.
- [5] (a) Find the solution set y = x 3,  $x^2 + y^2 = 17$

- (b) A card is drawn randomly from 30 identical cards numbered from 1 tp 30, find that the probability that the number on the drawn card is
  - (1) divisible by 4
  - (2) prime number

## Model (2)

## [1]Complete the following:

(1) if x = 2, y = 3, then  $(y - 2x)^{10} = \dots$ 

(2) If  $x \in R$ -  $\{0,3\}$ , then  $\frac{x}{x-3} \div \frac{x}{3-x} = \dots$  in its simplest form

(3) If A ,B are two events in a random experiment, and  $A \subset B$ , then  $P(A \cup B) = ...$ 

(4) If x - y = 3, x + y = 9, then y = ...

(5) the common domain of two fractions  $\frac{x}{x^2-1}$ ,  $\frac{3}{x^2+x}$  is ......

(6) If  $\{-2, 2\}$  is the set zeros of the function  $f(x) = x^2 + a$ , then  $a = \dots$ 

## [2]Choose the correct answer from the given ones

(1) If the sum of two numbers is 8, and their product is 15, then the two numbers are ......

(a) 2,6

(b) 3.5

(c) 4.4

(d) 1,15

(2) If the fraction  $\frac{x-a}{x+5}$  is the multiplication inverse of  $\frac{x+5}{x+3}$ , then  $a = \dots$ 

(a) 3

(b) - 5

(c) - 3

(d) 5

(3) If a coin is tossed once, the probability that the head appears = ......

(a) 1

(b)  $\frac{3}{4}$ 

(c)  $\frac{1}{2}$ 

(d)  $\frac{1}{4}$ 

(4) If x + y = 0,  $x^2 = 25$ , then  $y = \dots$ 

(a) 20

(b) - 5

(c) 5

 $(d) \pm 5$ 

(6) If the probability that Hamdy succeeded is 95%, then	
probability that he	
does not succeed is	
(a) 20% (b) 10% (c) 5% (d) zero	0
(a) 20% (b) 10% (c) 5% (d) zero [3](a) solve the two equations: $x - 2y = 1$ , $x^2 - xy = 0$	
(b) Find n in the simplest form showing its domain where	9
$\mathbf{n}(\mathbf{x}) = \frac{\mathbf{x} - 3}{\mathbf{x}^2 - 7\mathbf{x} + 12} - \frac{4}{\mathbf{x}^2 - 4\mathbf{x}}$	
[4](a) solve the equation $2x^2 - 5x + 1$ approximate to the	
nearest two decimals	
(b) If $n(x) = \frac{x^2 - 3x + 2}{x^2 - 1} \div \frac{3x - 15}{x^2 - 4x - 5}$ Find n in the simplest form	
showing its domain.	
[5] (a) If the set of zeros of the function f where	
$f(x) = ax^2 + bx + 8$ is {2,4}, find the value of a and b	
(b) A ball drawn randomly from 25 identical balls of the same	•
volume and weight, 10 of them are red, 8 are white and the	
rest are green, find the	
probability that the drawn ball is :	

(5) If the two equations x + 3y = 6, 2x + ky = 12 have an

(c) 3

(d) 1

infinite number of

solutions, then  $k = \dots$ 

(a) 2 (b) 6

(1) White (2) Green or white (3) Not green

## Model (3)

## [1]Complete the following:

(1) If  $x \in \mathbb{R}$ - { 2 }, then  $\frac{x}{x-2} + \frac{x}{2-x} = \dots$  in its simplest form

(2) If 
$$x y = 3$$
,  $x^2y - xy^2 = 6$ , then  $x - y = \dots$ 

- (3) If A ,B are two mutually exclusive events in a random experiment, then  $P(A \cap B) = ...$
- (4) If the length of a rectangle is 3 cm. and its diagonal of length 4 cm. Then its width is ........... Cm

(5) If 
$$(5, x-4) = (y+2,3)$$
, then  $x + y = ....$ 

(6) The common domain of two fractions  $\frac{5}{x-3}$ ,  $\frac{x+1}{x^2+3x}$  is ......

## [2]Choose the correct answer from the given ones

(1) If x is an negative number, then the greatest number is ...

(a) 
$$7 + x$$

(c) 
$$7x$$
 (d)  $\frac{7}{x}$ 

(d) 
$$\frac{7}{x}$$

(2) If  $n(x) = \frac{x-1}{x+3}$ , then the domain of  $n^{-1}$  is .....

(a) 
$$R - \{ -3 \}$$

(b) 
$$R - \{1\}$$

(a) 
$$R - \{-3\}$$
 (b)  $R - \{1\}$  (c)  $R - \{1,-3\}$  (d)  $\{1,-3\}$ 

(3) If a die is tossed once, then the probability of appearance of an odd number equals

(a) 
$$\frac{1}{3}$$

(b) 
$$\frac{1}{2}$$

(4) If the solution set of the equation:  $x^2 - ax + 4 = 0$  is  $\{-2\}$ , then  $a = \dots$ 

- (a) zero
- (b) 1
- (c) 2 (d) 4

(5) If the two equations x + 2y = 4, 2x + ky = 11 represent two parallel lines

, then  $k = \dots$ 

- (a) 4
- (b) 4

- (d) 1

(6) If the solution set of the two equation: x - y = 0, xy = 16 is..

- (a)  $\{(0,0)\}$  (b)  $\{(4,4)\}$  (c)  $\{(-4,-4)\}$  (d)  $\{(4,4),\{(-4,-4)\}\}$

[3] (a) Find n in the simplest form showing its domain where

$$n(x) = \frac{x^3 - 8}{x^2 + x - 6} \times \frac{x + 3}{x^2 + 2x + 4}$$

(b) Find the solution set of the equation  $(x-3)^2 - 5x = 0$ approximate to the nearest two decimals.

[4] (a) If  $n(x) = \frac{x+5}{x^2+7x+10} - \frac{x-1}{x^2+5x+6}$ , find n in the simplest form showing its domain, find n(-2)

(b) Find the solution set of the two equations: x - y = 4, 3x + 2y = 7 Graphically then verify algebraically.

[5] (a) Graph the function f where  $f(x) = 4x - x^2 - 3$  on the interval [0,4]. Find the

## Unit 4 Lesson 1 : Basics on circle

## 1- Complete the following:

- 1. If one end of a line segment lies on the center of the circle and the other end on the circle, then this line segment is called ........
- 2. If the two ends of a line segment lie on the circle, then this line segment is called ......
- 3. The chord which passes through the center of the circle is called ......
- 4. The straight line which passes through the center of the circle is called ......
- 5. The longest chord of the circle is called .......
- 6. The circle has ...... Number of axes of symmetry.
- 7. In any circle the perpendicular straight line on any chord its mid point is ...... to the circle
- 8. The circle divides the plane into ...... sets of points

## 2- In the opposite figure:

 $\overline{AB}$  is a diameter of a circle M ,  $\overline{AC}$  is a chord  $\overline{MD} \perp \overline{AC}$  , m ( $\angle A$ ) = 30°**Prove that**:(1)  $\overline{MD}$  //  $\overline{BC}$  is an equilateral triangle

3- In the opposite figure: $\overline{AB}$ and $\overline{BC}$ are two chords in circle M  Which has radius length of 5 cm, $\overline{MD} \perp \overline{AB}$ intersects $\overline{AB}$ At D and intersects the circle M at E,
X is the midpoint of $\overline{BC}$ , AB = 8 cm, m( $\angle ABC$ ) = 56° Find: 1) m ( $\angle DMX$ ) 2)length of $\overline{DE}$
4- In the opposite figure :  M is a circle of $r = 13$ cm $\sqrt{AB}$ is a chord of length 24 cm
C is the midpoint of $\overline{AB}$ and $\overline{MC} \cap circle\ M = \{D\}$ Find: The area of $\triangle$ ADB
5- In the opposite figure  AC = AB , X is the midpoint of $\overline{AB}$ ,  Y is the midpoint of $\overline{AC}$ , m( $\angle MXY$ ) = 30°  Prove that : The triangle AXY is equilateral $\triangle$
6) In the opposite figure: Prove that: The triangle XYM is an equilateral $\triangle$
C $E$ $120$ $A$ $B$

7)In the opposite figure: In circle M, $\overline{MX} \perp \overline{AB}$ , $\overline{MY} \perp \overline{AC}$ $m(\angle A) = 60^{\circ}$ and $m(\angle B) = 70^{\circ}$ Find: The measures of the interior angles Of the triangle MXY	C $A$
8) In the opposite figure: $\overline{MD} \perp \overline{BC}$ and $\overline{ME} \perp \overline{AC}$ Prove that:  1) $\overline{ED}$ // $\overline{AB}$ 2) The perimeter of $\Delta$ CDE = $\frac{1}{2}$ the perimeter of $\Delta$ ABC	A C D B
9) In the opposite figure: $\overline{AB}$ is a chord of circle M , $\overline{AC}$ bisects $\angle$ BAM And intersects circle M at C If D is the midpoint of $\overline{AB}$ Prove that : $\overline{DM} \perp \overline{CM}$	A A D B

## Unit 4 Lesson2: Position of a point & straight line with respect to circle

## 1)Complete:

- 1) The St. line which is drawn  $\perp$  to a diameter of a circle at one of its endpoints is ......
- 2) The two tangents to a circle at the two endpoints of a diameter of it are......
- 3) If M is a circle with circumference  $8\,\pi$  cm , A is a point on the circle , then MA = ...... cm
- 4) If the St. L is a tangent to the circle whose length of diameter is 8 cm, then the St. is at a distance ...... cm from its centre.
- 5) M is a circle of radius length 5 cm A is a point on the circle where MA =  $(2 \times + 1)$  cm then x = ....
- 6) If the area of the circle M is  $16~\pi~cm^2$  , A is a point in its plane where MA = 8 cm , then A lies ...... The circle M
- 7)M is a circle with diameter of length 8 cm If the St. line L is outside the circle , then the distance between the centre of the circle and the St. line L  $\in$  ......
- 8)If M is a circle with radius length 7 cm ,  $\overline{MA} \perp L$  where  $A \in L$  , if the St. line L intersects the circle M and MA =  $3 \times -5$  , then  $\times \in ...$  And if the St. line L is a tangent to the circle M and MA =  $\times^2 2$  , then  $\times \in ...$
- 9)  $\overrightarrow{CA}$  touches the circle M at A , m ( $\angle ACM$ ) = 30° If the radius length of the circle M = 4 cm , then MC = ......... Cm

30

2- In the opposite figure: $\overrightarrow{BC}$ is a tangent at B, m ( $\angle C$ ) = 45°, D is the midpoint of $\overrightarrow{AH}$ Prove that: DA = DM
3- In the opposite figure:  M is a circle with radius length 5 cm, $XY = 12 \text{ cm}$ , $\overline{MY} \cap circle M = \{Z\}$ And $ZY = 8 \text{ cm}$ Prove that: $\overline{XY}$ is a tangent to the circle M at X
4- $\overline{AB}$ is a diameter in a circle of area $36\pi$ cm² , $\overline{BC}$ is drawn a tangent to the circle at B , if m ( $\angle$ $ACB$ ) = $60^\circ$ , then calculate the area of $\Delta$ ABC ?
5- In the opposite figure : Find the length of $\overline{BC}$ , B C = 44 cm, $\pi = \frac{22}{7}$ Find Area & circumference of the circle ?

6)Choose	the o	correct	answer:

1) If the point A is in the plane of the circle M whose radius length = r and  $\,$  if  $\,$  0 < MA < r  $\,$ , then A lies ......

[ on , inside , outside , axis of symmetry ]

2) A circle with diameter length 2 x cm , the St. line is at a distance (x + 1)cm from its center , then the St . line L is.....

[ tangent , secant , outside , axis of symmetry ]

3) If the diameter length of a circle is 8 cm and the St. line L is at distance of 3 cm from its center , then the St. line L is......

[ tangent , secant , outside , axis of symmetry ]


4)A circle is of a circumference 6  $\pi$  cm , and the St. L is distant from its center by 3 cm , then the St. line L is .......

[ tangent , secant , outside , axis of symmetry

## 7)In the opposite figure :


M is a circle  $\overrightarrow{AB}$  is a tangent to the circle at X MA = 8 cm, m (  $\angle$  ABM) = 30° and  $\overline{AC} \perp \overline{MB}$ 

Find: The length of each of  $\overline{AB}$  and  $\overline{AC}$ 



## 8)In the opposite figure :

Find the  $m(\angle ECM)$ 



## Unit 4 Lesson3: Position of a circle with respect to another circle

<u>1</u> )	Con	<u>npl</u>	<u>ete</u>

1) If the surface of the circle M $\cap$ the surface of the circle N = $\emptyset$ , then the two circles M and N are
2) If the surface of the circle M $\cap$ the surface of the circle N = { A } , then the two circles M and N are
3) The centres line of two intersecting circle is perpendicular to and
4 )The centres line of two Touching circles is perpendicular to
5) M and N are two circles touching internally, their radii lengths are 7 cm and 5 cm then MN = cm
6) M and N are two circles of radii lengths are 9 cm and 4 cm respectively, MN = 5 cm, then the two circles are
7) M and N are two circles, their radii lengths are 8 cm and 3 cm, if MN = 11 cm, then the two circles M and N are
- 7 cm, then the two circles want with and with a second
2- M and N are two intersecting circles at A and B , MA = 12 cm , NA = 9 cm and MN = 15 cm . Find : The length of $\overline{AB}$ ?

3- In the opposite figure:  M and N are two intersecting circles at A and B, C is the midpoint of $\overline{XY}$ , m( $\angle$ D) = 40°, FZ is a tangent to the circle N at F where $\overline{MN} \cap \overline{FZ} = \{F\}$ 1) Find: m( $\angle$ CME) 2) Prove that: $\overline{FZ}$ // $\overline{AB}$
4- In the opposite figure:  M and N are two intersecting circles at A and B, $C \in \overrightarrow{BA}$ , $D \in \text{the circle N}$ , $m (\angle MND) = 125^{\circ}$ $m (\angle BCD) = 55^{\circ}$ ,  Prove that: $\overrightarrow{CD}$ is a tangent to circle N at D
5- In the opposite figure:  M and N are circles with radii lengths of 10 cm  And 6 cm respectively and they are touching internally at A, $\overrightarrow{AB}$ is a common tangent for both . If the area of $\Delta$ BMN = 24 cm <sup>2</sup> , Find: The length of $\overline{AB}$

## 6)Complete:

1)If the radius length of the circle M = 3 cm and the radius length of the circle N = 5 cm , MN = 6 cm , then the two circles M and N are


2)M and N are two intersecting circles their radii lengths are 3 cm and 5 cm respectively , Then MN∈ ......

3)If the radius length of the circle M = the radius length of the circle N = MN, then the two circles are .....


3)M , N and L are 3 circles touching externally two – by – two , their radii length are 5 cm , 6 cm and 4 cm , then the perimeter of the triangle MNL = ......

4)M is a circle of radius length 4 cm , touches the circle N externally . If MN = 7 cm , then the circumference of the circle N = ......  $\pi$  cm

## 7) In the opposite figures:



$$AB = \cdots cm$$
.



## Unit 4 Lesson4: Identify the circle

1) It is possible to draw passing through a given point.
2) The number of circles which passes through two given points is
3) The number of circles which passes through three collinear points is
4) The number of circles passing through three non-collinear points is
5)The center of the circumcircle of a triangle is the point of intersection of
6)If $\Delta ABC$ is right – angled at B , then the center of its circumcircle is
7)It is impossible to draw a circle passing through the vertices of
2)Use the graph copy book to draw:  1)Draw $\overline{AB}$ with length = 4 cm, then draw a circle passing through the
Points A and B with radius length = 3 cm .How many solutions can be obtained?
2)Draw $\overline{AB}$ with length = 6 cm , then draw a circle passing through the two
Points A and B with radius length = 4 cm .How many solutions can be obtained?

### 3- Complete:

- 1- If the number of circles can be drawn passing through two given points in the plane equals ......
- 2- If two circles have three common points, then they are .......
- 3- The radius of the smallest circle drawn to pass through two given points in the plane equals ......
- 4- The point of intersecting of the symmetric axes of the sides of a triangle is ......
- 5- If M is a circle of radius r, A is a point in the plane of the circle:
  - (a) If  $MA = \frac{1}{2}R$  then A ..... the circle
  - (b) If MA = R then A ..... the circle
  - (c) If MA = 3R then A ..... the circle

## 4) Use the graph copy book to draw:

1)Draw the right – angled triangle ABC at B where AB = 4 cm , BC = 3 cm , then draw the circumcircle of this triangle . Where does the center of the circle lie

With respect to the sides of this triangle?

- 2)Draw the isosceles triangle ABC in which m( $\angle$  ABC) = 120°, BC = 4 cm. Determine the center of the circumcircle of it and find its radius length.
- 3)Draw the equilateral triangle ABC of side length of 6 cm ,Draw the circumcircle of this triangle ABC and located the position of the center of the circle .

## Unit 4 Lesson5: Relation between chords & centre of circle

1- <u>In the opposite figure :</u>	C
$\overline{AB}$ and $\overline{AC}$ are two chords equal in length in the circle M,	E
m ( $\angle$ CAB) = 70°, X is the midpoint of $\overline{AB}$ ,	Mry
Y is the midpoint of $\overline{AC}$	700
Find: 1) m (∠ DME)	BXA
2)Prove that: XD = YE	
2)In the opposite figure: $\Delta$ ABC is inscribed in the circle M  If DM = EM , m ( $\angle$ DME) = 120°  Prove that: $\Delta$ ABC is an equilateral $\Delta$	C E 120 D B
3) In the opposite figure $AB = AC$ , $D$ and $E$ are the midpoints of $\overline{AB}$ and $\overline{AC}$ respectively, $m$ ( $\triangle MDE$ ) = 30°  Prove that:  MDE is an isosceles $\triangle$ 2) ADE is an equilateral $\triangle$	A 30° B
4)In the opposite figure : Provethat: CD=XY	X C
	Y M D

5) In the opposite figure
ABC is a triangle in a circle of center M,
$\overline{MD} \perp \overline{AC}$ , $\overline{ME} \perp \overline{AB}$ and
BC = 8 cm Find DH
6) In the opposite figure
M circle of center M, its radius of length 13 cm
AB is a chord of length 24 cm,
C is the mid point
Of AB, MC cuts the circle at D. Find:
[1] The length of MC [2] the area of $\triangle$ ADB
7) In the opposite figure
AB and AC are two tangent to the circle M,
Touch it at B, C respectively
and m(∠BAM)=25°
[1] prove that $\overrightarrow{MA}$ bisects ( $\angle BMC$ ) [2] find ( $\angle BMC$ )
••••••

## **General Exercise on Unit four**

1- choose the correct from	the given ones:
1) If the length of a diameter at distant 3.5 cm from its cen	of a circle is 7 cm, and the straight line L tre , then L is
a) Secant to the circle at two	points b) Lies outside circle
c) Tangent to the	d) Axis of symmetry to the circle
2) If the point A belongs to the equals.	ne circle M of diameter 6 cm, then MA
(a) 3 cm (b) 4 cm	(c) 5 cm (d) 6cm
3) If the straight line L is a ta then the distance between L a	angent to the circle M of diameter 8 cm, and its centre equals
(a) 3 cm (b) 4 cm	(c) 6 cm (d) 8 cm
4) If the straight line L is out: M, if L at distance X from its	side a circle of radius 3 cm, and its center centre, then x∈
(a) ] 3, $\infty$ [ (b) [3, $\infty$ [	(c) ] 6, $\infty$ [ (d)]- $\infty$ , -6 [
5) If the straight line L at dist radius R,	ance X from a circle of centre M and
$X \in ]0, R[, then L$	
a) Intersects the circle	b) Touches the circle
c) Lies outside the circle d	) Passes through the centre of the circle
	dicular drawn from the centre of the circle 6 cm and the radius 6 cm, then L
a) Intersects the circle	b) Touches the circle
c) Lies outside the circle	d) Passes through the centre of the circle

	f the following poi e origin and its ra		long to the centre that its	S
a) (0,7)	b) (0, -7	c) (7,0)	<b>d</b> ) (7,7)	
8) If the sur	face of the circle	M∩ the surface	of the circle N={A}, the	n
the two circ	eles M and N are.	•••••		
a) Distant		b) Co	oncentric	
c) Touching	g externally	d) In	tersecting	
9) The num	ber of circles can	be drawn to pa	ss through the end point	ts
	egment AB equal			
a) 1	b) 2 c)	3 d)an infin	ite number	
10) If the ci	rcle M∪ the circle	$\mathbf{N} = \{\mathbf{A}, \mathbf{B}\}, \text{ then}$	the two circles M and N	J
are				
a) Distant		b) C	Concentric	
c) Touchin	g externally	d) In	tersecting	
11) If the ty	vo circles M,N are	touching exter	nally, the radius of one o	of
		_	of the other circle equals	
•••••				
(a) 3 cm	(b) 4 cm	(c) 7 cm	(d) 14 cm	

12) If the two circles M,N are touching internally, the radius of one of them 3 cm, and $MN = 8$ cm, then the radius of the other circle equals			
(a) 5 cm	(b) 6 cm	(c) 11 cm	(d) 12 cm
13) M and N then MN=		ing circles thei	r radius are 5 cm, 2cm,
(a) ] 3, 7[	(b) [3,7[	(c) ] 3, 7]	(d)[3,7]
equals	ber of circles that  o) One (c) Three		three collinear points nfinite number
=	metric axis of the circles M , N is		AB to the two
(a) MA	(b) MB	e)MN	(d) $\overrightarrow{NA}$
16) the centr	es of the circles w	hich pass thro	ugh the two points A , B lie
a) The axis	of AB	b) AI	3
c) The perpe	endicular to $\overline{AB}$	d) The	perpendicular on $\overline{AB}$ at B
17) Number equals		n pass through	three non collinear points
(a) zero	(b) one	(c) two	(d)three

c) its height			
d) The symmetric axis of its sides			
19) If the two points A, B lie on a plane AB= 4cm, then the length of the radius of the smallest circle passes through A and B equals			
(a) 2 cm (b) 3 cm	(c) 4 cm (d) 8 cm		
20) If the two points A, B lie on a plane AB= 6cm, then the number of circles of each of them has a radius of 5 cm and passes through A and B equals  a) zero b) 1 c) 2 d) an infinite number  2: Match from the column (X) to the column (Y) to get a true statement  Two circles of radius cm. & 6 cm			
X	Y		
1) If $MN = 1$ cm	a) M, N are two intersecting circles		
2) If MN = 2 cm	b) M , N are two distant circles		
3) If MN = 7 cm	c) M, N are two touching externally		
4) If MN = 14 cm	d) M, N are two interior circles		
5) If MN = 15cm	e) M, N are two touching internally		
17			

18) The centre of the circum circle of any triangle is the point of

intersecting of

a) Interior bisectors of its angles

b) Exterior bisectors of its angles

its .....

# 3- in the opposite figure A circle of circumference 44 cm, $\overline{AB}$ is a diameter $\overrightarrow{BC}$ is a tangent at B , and m ( $\angle C$ ) = $60^{\circ}$ Find the length of $\overline{BC}$ ( $\pi = \frac{22}{7}$ ) 4- In the opposite figure $\overline{AD}$ is a diameter in a circle M, $\overrightarrow{CA}$ and $\overrightarrow{CB}$ are two tangent to the circle M, touch it at B, C respectively. Prove that $m (\angle DMB) = m (\angle ACB)$


5- In the opposite figure


M, N are two intersecting circles,

MN intersects the circle

M at C,  $\overline{\text{CA}}$  is a tangent to the circle M at C,

and cuts the Circle N at A, B Prove that:






6- In the opposite figure

M, N are two intersecting circle,

 $\overline{CD}$  is a chord in the circle M, cuts  $\overrightarrow{MN}$  at E, if E is the mid point

of  $\overline{CD}$  Prove that:  $\overline{AB}$  //  $\overline{CD}$ 




7) M, N are two touching internally circles at A, the circle M is greater than circle N, draw the common tangent  $\overrightarrow{AC}$ , then draw  $\overrightarrow{NM}$  to cut the circle N at B, and draw the tangent  $\overrightarrow{BD}$  to the circle N to cut the circle M at D, E Prove that: [1]  $\overrightarrow{AC}$  //  $\overrightarrow{BD}$ [2] BD =BE. 8) In the opposite figure M, N are two congruent circles,  $\overline{AC}$  is a tangent to the circle M at A the,  $\overline{DF}$  is a tangent to the circle N at D,  $\overline{AC}$  //  $\overline{DF}$  . Prove that: [2] AB = ED[1] BC = FE

## 9) In the opposite figure

 $\overline{AB}$  ,  $\overline{CD}$  are two chords (equal in length) if X,Y are the two mid points of  $\overline{AB}$  ,  $\overline{CD}$ 


respectively Prove that



[1] 
$$\overline{AC}$$
 //  $\overline{BD}$  [2] m ( $\angle BXY$ ) = m ( $\angle DYX$ )


## 10) In the opposite figure

M, N are two intersecting circles at A, B,



X is the mid point of  $\overline{AC}$ 

Prove that: NX = NY.




11) $\overline{AB}$ , $\overline{CD}$ are two parallel chords in a circle M, E is the midpoint of $\overline{AB}$ , $\overrightarrow{EM}$ is drawn to cut $\overline{CD}$ at F.
Prove that: FC = FD
12) A, B are two points where AB = 6 cm, Draw a circle of radius 5 cm and passes through the two points A, B. Find the distance from the centre to $\overline{AB}$ .
13) Draw the triangle ABC in which AB = 6 cm, AC = 4 cm, m ( $\angle BAC$ ) =60°.
Then draw a circle passes through the two points A, C and its center $\in \overline{AB}$ .

14)  $\overline{AB}$  is a diameter in a circle M,  $\overline{AC}$  is a chord such that m ( $\angle BAC$ ) =  $30^{\circ}$ , then draw ,  $\overline{BC}$  and ,  $\overline{MD} \perp \overline{AC}$  to cut it at D. Prove that: [1]  $\overline{MD}$  //  $\overline{BC}$  [2] BC = the length of the radius of this circle.


#### Unit 5 **Lesson 1 : Central angles**

### 1)Complete:

- 1)If a square ABCD is inscribed in a circle M, then m(AB) = .....
- 2) The circumference of a circle = 36 cm , then the measure of an arc of it with length = 6 cm is ......
- 3)An arc in a circle, its length =  $\frac{1}{3}\pi$  r, then it is opposite to a central angle of measure ......
- 4) The measure of the arc which represents  $\frac{1}{3}$  the circumference of the circle =
- 5)In the opposite figure: m(AC) =




## 2)In the opposite figure Find: $m (\angle ABC)$




## 3)In the opposite figure:

Find the length of :  $\widehat{AB}$  (  $\pi = \frac{22}{7}$ )




## 4)In the opposite figure:

Find : m (BE)		




## 5)In the opposite figure:

Prove that : m (	




## 6)In the opposite figure:

Prove that: $m(\widehat{AB}) = m(\widehat{AD})$



## 7)In the opposite figure:


Prove that :∆ MCD is equilateral	



## 8)In the opposite figure:

AB is a diameter of circle M m (AC): m (DB) = 5:6


Find : m ( ACD )	



## 9)In the opposite figure :

 $\overrightarrow{CD}$  is a tangent to the circle at C

Where  $\overline{AB}$  //  $\overline{EF}$  //  $\overline{CD}$ Prove that: CE = CF




# Unit 5 Lesson 2 :Relation between central angles & inscribed angles with same arcs


## 1)In the opposite figure:

M is a circle. In each of the following, find the value of the symbol used in measuring:


1




2



3




4




z = .....

5



 $\chi = \cdots \cdots$ 

6




y = ...... z = .......

2)In the opposite figure:

Find:  $m (\angle BCD)$ ,  $m (\angle BAD)$ 

.....



3)In the opposite figure :	A
Find :the area of the circle M ( $\pi = \frac{22}{7}$ )	45 M
	14\2cm.
	14420
	C
4)In the opposite figure :	C
<b>Find</b> : m ( ∠ <i>MBC</i> )	B
	120 A
	130
	1
5)In the opposite figure :	, n
Prove that: m ( $\angle AMC$ ) = m ( $\angle ADB$ )	D
	/ / M \ \
	B A
6)In the opposite figure :	
Prove that : BE> AE	
	/ M
	M
	\ E


## Unit 5 Lesson 3: Theorem 1 & corollaries

## 1)Complete:

- 1) The measure of the arc that is opposite the inscribed angle of measure  $50^{\circ}$  = ...
- 2) The inscribed angle which is drawn in a semicircle is.......
- 3) If two chords intersect at a point inside a circle, then the measure of the included angle equal ...... of the sum of the two measures of the two arcs.
- 5) The measure of the inscribed angle which is drawn in  $\frac{1}{3}$  a circle equals .......


## 2)In the opposite figure:

 $\overline{AB}$  is a diameter in circle M , m ( $\angle$ C) = 20° , m ( $\overline{BE}$ ) = 70° Then m ( $\overline{DE}$ ) = ......




## 3)In the opposite figure:


m (CE) = .....°



## 4 Complete the following:



If 
$$m(\widehat{AC}) + m(\widehat{BD}) = 80^{\circ}$$
,  
then  $m(\angle AEC) = \dots^{\circ}$ 



If m (
$$\angle$$
 DEB) = 110°,  
m ( $\widehat{BC}$ ) = 70°, then m ( $\widehat{AD}$ ) = ......°

## 5 In the opposite figure:


XYZ is a triangle inscribed in the circle N,

$$m (\angle XYN) = 40^{\circ}, m (\angle NZY) = 30^{\circ}$$

## Complete:


$$3 \text{ m}(\widehat{XZ}) = \dots^{\circ}$$

4 m 
$$(\widehat{XY}) = \dots \circ$$




## 6)In the opposite figure:

$$m(AC) - m(BD) = 70^{\circ}$$
  
then  $m(\angle E) = \dots$ 




## 7) In the opposite figure:

 $\overline{AB}$  is a diameter in circle M , r = 4 cm , m ( $\angle$  A) = 30 ° , BC = ...... cm




# 8)In the opposite figure:

If m (BC) = $112^{\circ}$	, m ( DE ) = $44^{\circ}$ ,	DA = AE
Then $m (\angle ADE)$	=	




# 9)In the opposite figure:

$\overline{AB}$ is a diameter in the cire	cle M
$m(\angle BMD) = 50^{\circ}$ Find with	h proof: $m(\angle ACD)$




10)In the opposite figure: Find with proof: m(∠CBD)



11)In the opposite figure:

Find with proof:  $m(\angle DCA)$ ,  $m(\angle CAB)$ 




# Unit 5 Lesson 4: Inscribed angles with

1)In the opposite figure :	$\bigcap$ D
Find: m (∠DEB)	
	n 125°
	$B \stackrel{125}{\longrightarrow} A$
	E
2)In the opposite figure:  Prove that: $m (\angle AEB) = m (\angle AEC)$	C
	E A
	B
3)In the opposite figure:	C
Find: m (∠AED)	
	B (55°
	MA
	E
4)In the opposite figure:	C A
Prove that: AF = FB	$\mathcal{K}$
	/   \ <sub>F</sub> /   \
	( A X A )
	<i>X V</i>
	$\mathcal{L}$


5)In the opposite figure:  Prove that: m (∠ DAC) = m (∠ BAE)	E B
6)In the opposite figure : Find with proof : m (BD)	B 30 A
7)In the opposite figure: Find:m(∠ACB)	C A B

# 8)In the opposite figure:

m(BD) = 6 Find with p	0°, m(AD) = 1 proof: m(CB)	100°, m ( <b>, m (</b> .	$\widehat{AC}$ ) = 120° $\angle ACB$ )	,



9)In the opposite figure: If m ( $\angle$ A) = 40°, m (BD) = 60° Find with proof: m (CB), m (EC)




#### Unit 5 **Lesson 5:** Cyclic quad & it's prop.


& theorem 3 with its converse same arc

# 1)In the opposite figure:

m (AB) =  $110^{\circ}$  and m( $\angle$ CBE) =  $85^{\circ}$ 

Find: m (∠BDC)






# 2)In the opposite figure:

Prove that: The figure ABDE is a cyclic quadrilateral


 $m(\angle CED) = \frac{1}{2} m(\widehat{AC})$ 



3)In the opposite figure:

Prove that: AD = DC





# 4)In the opposite figure:

 $m (\angle BDC) = 90^{\circ}$ ,  $m (\angle ABC) = 60^{\circ}$ 

$$m (\angle ACB) = 30^{\circ}$$

Prove that: The figure ABCD is a cyclic quadrilateral determine where is the center of the circle passes through the vertices of figure ABCD

5)In the opposite figure: Prove that : AXEY is a cyclic quadrilateral	A $C$ $A$ $X$ $E$ $X$ $B$
6)In the opposite figure: Prove that: 1)The points D, E, C and B have one circle past Through them. 2) m(∠AME) = 2 m (∠ D)	sing D  B  C  M
7)In the opposite figure:  Prove that: 1)The figure AMEC is a cyclic quad 2)m(∠C)	rilateral D A

8)In the opposite figure:	<u>.</u>
Prove that: 1)The figure AEFD is a cyclic quadrilate	eral
$(2)\overline{EF} //\overline{BC}$	
	A
DZ	
	\ /*\\\
	$\times$ /
	XXE /
	1 /
C	В
9)In the opposite figure:	X
Prove that: The figure AFXE is a cyclic quadrilatera	al 🖊
	E/ /
	AE
	/()   )
	/ ( ) ( )
X	
	$\mathcal{L}_{B}$
10)	
10)In the opposite figure:	

m ( $\angle$  C) = 50°, m ( $\angle$  A) = 80° AD = AB ,**Prove that: The points A , B , C and D have one** Circle passing through them.


# Unit 5 Lesson 6: Relation between tangents of circle

1)In the opposite figure :
Prove that : AB = CD
$\begin{pmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \mathbf{i} & \mathbf{j} & \mathbf{k} \end{pmatrix} \begin{pmatrix} \mathbf{i} & \mathbf{k} \\ \mathbf{k} & \mathbf{k} \end{pmatrix}$
$\dot{N}$ $\dot{R}$ $\dot{M}$ $\dot{M}$ $\dot{M}$
fig. (1) fig. (2)
2)In the opposite figure:
Prove that:
1) $\overrightarrow{AB}$ bisects $\angle DAX$
$(2)\overline{AD} //\overline{XB}$
X 470°
A
3)In the opposite figure:
Prove that:
1) $\overrightarrow{CB}$ bisects $\angle ACD$
2) Find: $m(\angle A)$
В
4)In the opposite figure :
Prove that:
1) C is the midpoint of $\overline{AB}$
$2)\overline{AD} \perp \overline{BD}$
•


# 5)Complete:

- 1) The two tangents drawn to the circle at the two ends of a diameter in it are ......
- 2) The two tangent segments drawn to a circle from a point outside it are ......
- 3) The number of common tangents of two distant circle is ......

4)In the opposite figure:	
$m ( \angle X ) = \dots$	




5)In the opposite	e figure :	
$m (\angle CAB) =$		



6)In the opposite figure:

m (∠CAB) =	




# 7)In the opposite figure:

m ( ∠ C ) = .....

A B


# 8)In the opposite figure:

Perimeter of Δ ABC = .....



# 9)In the opposite figure:

m ( ∠ CBM ) = .....



#### Unit 5 **Lesson 7:** Angles of tangency& converse of theorem 5


1)	Com	nlat	Δ.
$\perp$	COIII	piei	<u>.e.</u>

1)The	angle of tangency is the included angle	e between ,

- 2) The measure of the angle of tangency equals the measure of .....subtended by the same arc .
- 3) The measure of the angle of tangency equals half the measure of .....subtended by the same arc


# 4) In the opposite figure:

$$m(\angle ADB) = \dots$$



# 5)In the opposite figure:


$$m(\overrightarrow{BC}) = \dots, m(\overrightarrow{AC}) = \dots$$
 $m(\angle ACD) = \dots$ 



# 2)In the opposite figure:

Find the value of x





3)In the opposite figure : Find m(∠A)  125  B
4)In the opposite figure: Prove that:  1)The figure ABCD is a cyclic quad.  2) \$\overline{AE}\$ is a tangent to the circumcircle of the quad. ABCD \$\overline{ABCD}\$ \$\overline
5) In the opposite figure : Prove that : 1) $\overline{BE}$ // $\overline{AC}$
2) Find: $m(\angle CDE)$ 3) $\overline{CE}$ is a tangent segment to the circle Passing through the points A, B, C
6)In the opposite figure : Prove that :
$\overrightarrow{AC}$ is a tangent to the circumcircle of $\triangle$ ABD

7)In the opposite figure :	
Find m(BC)	C B A X
8) In the opposite figure : Find m(BC)	A 60° M
9)In the opposite figure: Find : m(∠CDB)	D A A 40 X
10)In the opposite figure: Prove that: 1) ZE = ZY 2) XZ // YE	Z D 130° E

# **General Exercise on Unit five**


#### First: Complete the following

- (1) The two opposite angles in the cyclic quadrilateral are ...........
- (2) The chords which opposite to equal arcs in any circle are ..........
- (3) The measure of the inscribed angle equals half the measure of .......
- (4) In the opposite figure

In a circle M, m ( $\angle A$ ) =  $48^{\circ}$ , then:

[1] m (
$$\angle C$$
) = ......

[2] m 
$$(\widehat{BD})$$
 = ......... " $(\widehat{BD})$  is the major arc'




- (5) The quadrilateral is said to be a cyclic quad. If the measure of an exterior angle at any vertex equals the ....... of the angle which opposite to its adjacent.
- (6) In opposite figure

In a circle M, m ( $\angle CAE$ ) = 36°, then:

(a) m (
$$\angle EBC$$
) =.....

(b) m (
$$\angle EMC$$
) = ......

(c) m (
$$\angle EDC$$
) = .....





- (7) The inscribed angle which opposite to a major arc in a circle is ........
- (8) The two parallel chords in a circle intercept two ...... arcs
- (9) The measure of an arc of a circle equals double .........
- (10) The two inscribed angles subtended in the same arc in a circle are ......
- (11) The altitudes of any triangle are .......


(12) In the opposite figure


In a circle M,  $\overline{AB}$  is a diameter,

 $\overrightarrow{CD}$  is a tangent at D, m ( $\angle BAD$ ) =  $28^{\circ}$  , then:









(14) The measure of the angle of tangency equals .... the central angle on its common arc.

(15) The number of all common tangent drawn to two distant circles equals ......

(16) The centre of the inscribed circle of any triangle is the point of intersection of

Second: Choose the correct answer from the given ones

(1) The two opposite angles in the cyclic quadrilateral are ......

(a) equal

(b) complementary

(c) supplementary

(d) alternate

(2) The center of the inscribed circle of any triangle is the point of intersection of its

(a) altitudes

(b) medians

(c) bisectors of its angles

(d) axes of symmetry of its sides

(3) The inscribed angle which opposite to the minor arc in a circle is ......

(a) reflex

(b) right

(c) obtuse


(d) acute

(4) In the opposite figure

In circle M,m ( $\angle AMB$ ) = 52°, then

 $m(\angle ABD) = \dots^{\circ}$ 

- (a) 52°
- (b) 104°
- (c) 128°




(5) In the opposite figure

 $\overline{AC}$ ,  $\overline{BD}$  are two intersecting chords in a circle M,

If  $m(\angle A) = 35^{\circ}$  and  $m(\angle AED) = 115^{\circ}$ ,

then m( $\widehat{AD}$ )= ......°

- (a) 70°
- (b)  $80^{\circ}$
- (c) 115°
- (d) 160°



115°

(6) In the opposite figure

 $\overline{AB}$  is a diameter in a circle M,

 $m(\angle ABC) = 40^{\circ}$ , then  $m(\widehat{BC}) = \dots$ 

- (a) 40°
- (b) 50°
- (c) 90°
- (d) 100°


(7) In the opposite figure

 $\overline{AB}$  is a diameter in a circle M,

 $m(\angle ABD) = 25^{\circ}$ , then



- (a) 25°
- (b) 50°
- (c)  $65^{\circ}$



C


- [2]  $m(\angle DCB) = \dots$ 
  - (a) 50°
- (b)  $100^{\circ}$
- (c) 115°
- (d) 125°

(d) 90°

(8) In the opposite figure

Two concentric circle at M,  $\overline{AB} \cap \overline{CD} = \{M\}$ ,

If  $m(\widehat{BD})$  = 80°, then  $m(\widehat{AC})$ =.....




#### Third: Answer the following question

(1) (a) Prove that the two opposite angles in a cyclic quad are supplementary.

#### (b) In the opposite figure

ABCD is a quadrilateral in which AB = AD ,  $m(\angle ABD) = 30^{\circ} \text{ and } m(\angle C) = 60^{\circ} \text{Prove that ABCD is a cyclic quad}$ 



(2) ABCD is a cyclic quadrilateral in which  $\overline{AB}$  //  $\overline{CD}$ , if E is the mid point of  $\widehat{AB}$  . Prove that EC = ED ?


# 3) In the opposite figure

 $\overrightarrow{MC} \cap \overrightarrow{AB} = \{c\}$ ,  $\overline{MC}$  intersect the circle a D.

 $m(\angle MAB)=20^{\circ}$ , Find

[1]  $m(\widehat{AD})$ 

[2] m(∠*DEB*)



- (4) ABC is an acute angled triangle drawn inside a circle. Draw  $\overrightarrow{AD} \perp \overline{BC}$  to cut  $\overline{BC}$  at D and cuts the circle at E circle, then draw  $\overrightarrow{CN} \perp \overline{AB}$  to cut  $\overline{AB}$  at N. Prove that :
  - [1] ANDC is a cyclic quad
  - [2]  $m(\angle BND) = m(\angle BED)$

- (5) ABC is an equilateral triangle drawn inside a circle, D is a point on the arc  $\widehat{AB}$ , E is a point on  $\overline{DC}$  such that AD = DE. Prove that:
  - [1] ADE is an equilateral triangle

[2] 
$$\overline{DB}$$
 //  $\overline{AE}$ 

[3] 
$$m(\angle DCB) = m(\angle EAC)$$

$$[4] DE = EC$$

# (6) In the opposite figure

ABC is a triangle in which AB = AC .  $\overline{BC}$  is a chord

In the circle M, if  $\overline{AB}$  ,  $\overline{AC}$  cut the circle at D, E


Prove that:

$$\overline{BC}$$
 //  $\overline{DE}$ 

and if m( $\angle DCA$ )= 30° and m( $\angle A$ ) = 50°,



[1] 
$$m(\angle BEC)$$
 [2]  $m(\angle BMC)$  [3]  $m(\angle CDE)$ 



(7) (a) prove that the angles subtended by the same arcs in the circle are equal in measure.


#### (b) In the opposite figure

ABC is a triangle in a circle,  $\overline{BX} \perp \overline{AC}$ ,  $\overrightarrow{AY} \perp \overline{BC}$ 

Cuts it at Y and cuts the circle at Z,


Prove that:

- [1] ABXY is a cyclic quad
- [2]  $\overline{BC}$  bisects  $\angle XBZ$



# (8) In the opposite figure


 $\overline{AB}$  is a diameter of a circle M, C  $\in$  the circle,  $m(\angle CAB) = 30^{\circ}$ , D is the mid point of the arc  $\widehat{AC}$  and  $\overline{DB} \cap \overline{AC} = \{E\}$ .



- [1] Find  $m(\angle BDC)$ ,  $m(\angle ABD)$
- [2] Prove that  $\Delta$  ABE is an isosceles triangle.

# (9) In the opposite figure

 $\overline{AB}$  is a diameter of a circle M, D is the mid point of the arc  $\widehat{AC}$  Draw  $\overrightarrow{DM}$  to cut the circle at,  $\overrightarrow{BF}$  is a tangent to the circle to cut  $\overrightarrow{AC}$  at F.



Prove that:

[1] MBFD is a cyclic quad.

[2] 
$$\overline{DM}$$
 //  $\overline{BC}$ 

# (10) In the opposite figure


$$m(\angle BAC) = m(\angle BDC) = 90^{\circ}$$
.

E is the mid point of  $\overline{BC}$  and m( $\angle AED$ ) = 48°.

- [1] Find  $m(\angle ABD)$
- [2] Prove that:

(a) 
$$m(\angle ABD) = m(\angle ACD)$$

(b) 
$$m(\angle AEC) = 2m(\angle ABC)$$



# **Models Exams in Geometry**


# Model (1)

#### [1] Complete the following:

(1) The radius of the circle is the line segment joining between.....

#### (2) In the opposite figure

If 
$$(\widehat{AC}) = 100^{\circ}$$
, m  $(\widehat{BD}) = 60^{\circ}$ ,  
then m( $\angle BND$ ) = ......



(3) Any three points not belonging to a straight line then ..... Passes through them.


(4) In the same circle, the chord of the equal arcs in measure are.......

(5) The measure of the inscribed angle equals half the measure of ......

# (6) In the opposite figure

If M is a circle, m ( $\angle$ A) = 48°, then:

[2] m (
$$\widehat{BD}$$
) = ...... "major"



[2] Choose the	correct fro	m the given o	ones:	
(1) Axis of symr M and N is		e common ch	nord $\overline{AB}$ for two intersecting circle is	
(a) <i>MA</i>	(b) <i>MB</i>	(C) <i>MN</i>	(d) $\overrightarrow{NA}$	
cm . The position	on of with r	espect this ci		
(a) inside the circle (c) on the circle			) outside the circle ) coincide on the centre	
(3) If M a circle				
at distance 3 cr (a) touches the			secant to the circle	
(c) lies outside	the circle	(d) a	axis of symmetry to the circle	
(4) The two tan circle are	gent which	drawn from	the two ends of a diameter of a	
(a) parallel		(b) €	equal in length	
(c) congruent	(c) congruent (d) intersecting			

(5)  $\overrightarrow{AB}$  &  $\overrightarrow{AC}$  are two tangents, m ( $\angle A$ ) = 60°,

If 
$$\overline{AB}$$
 = 4cm, then  $\overline{CB}$  = ..... cm


- (a) 3
- (b) 4
- (c) 5
- (d) 8

# (6) In the opposite figure

M is a circle, m (
$$\angle$$
BCD) = 130°

Then m (∠BMD) = ......°

- (a) 50
- (b) 230
- (c) 100
- (d) 260




# (3) (a) In the opposite figure

Two concentric circles at M,  $\overline{AB}$  ,  $\overline{AC}$  are two

Tangents to the smaller circle m ( $\angle A$ ) = 70°.

- [1] Find m ( $\angle$ DME).
- [2] Prove that AB = AC.



Find the measure of all angles of the triangle ABC.

#### (4) In the opposite figure


 $\overline{AB}$  is a diameter of a circle M,

 $C \in \text{the circle, m } (\angle CAB) = 30^{\circ},$ 

D is midpoint of

$$\widehat{AC}$$
,  $\overline{DB} \cap \overline{AC} = \{E\}$ .

- [1] Find m ( $\angle$ BDC), m ( $\angle$ ABD)
- [2] Prove that:  $\triangle$  ABE is an isosceles triangle.



# (5) (a) State three cases of cyclic quadrilateral

- **(b)**  $\overline{AB}$ ,  $\overline{CD}$  are two orthogonal chords in a circle M, intersecting at E, draw  $\overrightarrow{BF} \perp \overrightarrow{AC}$  cuts it at  $\mathbf{F} \not\in \overline{AC}$ . Prove that:
  - [1] FCEB is cyclic quadrilateral
  - [2] m ( $\angle$ BFE) = m ( $\angle$ BAD)

# Model (2)


# {1} Complete the following:

- (1) The diameter passing through the mid point of any chord is ......
- (2) The number of symmetric axes of a circle equals .........
- (3) The line of centers of two touching circle is perpendicular to ........
- (4) The figure is said to be cyclic quad if the measure of any exterior angle at any vertex equals...... The opposite to its adjaçent.

# (5) In the opposite figure:

 $\overline{AB}$  is a diameter in a circle M,

If m 
$$(\widehat{AC})$$
 = m  $(\widehat{CD})$  = m  $(\widehat{DB})$ , then  
m  $(\angle CMD)$  = ......°, m  $(\angle CED)$  = ......



# [2] choose the correct from the given ones:


- (1) if  $\overline{AB}$  is a line segment, then the number of circles can be dawn passing through A, B equal ......
  - (a) 1
- (b) 2
- (C) 3
- (d) infinite number
- (2) If the straight line L  $\cap$  the circle M =  $\emptyset$ , the straight line L is ......
  - (a) outside the circle
- (b) secant to the circle
- (c) tangent to the circle
- (d) axis of symmetry of the circle

#### (3) In the opposite figure

If M is a circle, m ( $\angle$ M) – m ( $\angle$ A) = 50°,

Then m ( $\angle A$ ) = ......

- (a) 40°
- (b) 50°
- (c)  $100^{\circ}$
- (d)  $130^{\circ}$




#### (4) In the opposite figure

 $\overrightarrow{AB}$  is a tangent to circle M,

If 
$$MB = 5 \text{ cm}$$
,  $AC = 8 \text{ cm}$ , then  $AB = \dots \text{cm}$ 

- (a) 5cm
- (b) 10cm
- (c)12cm
- (d) 13cm



(5) All the centres of the circle which passes through

the two point A, B lie on......,

- (a) the axis of  $\overline{AB}$
- (b)  $\overline{AB}$
- (c) the perpendicular to  $\overline{AB}$
- (c) the mid point of  $\overline{AB}$
- (6) If the length of an arc of a circle is  $\frac{1}{3}\pi$  r, then its opposite central angle of measure .......
  - $(a) 30^{\circ}$
- (b) 60°
- (c) 120°
- (d) 240°
- (3) (a) Draw the circle which passes through the vertices of  $\Delta$  ABC such that AB = 3 cm , BC =4cm and AC = 5cm .

# (b) In the opposite figure

 $\overline{AB}$  is a tangent to circle M,


 $\overline{AC}$  is a secant, where E is the mid point of  $\overline{CB}$  ,

D is the mid point of  $\overline{AC}$ ,



Prove that

- [1]  $\overline{AB}$  //  $\overline{DE}$
- [2] The points N, D, C and E lie on one circle.



**(4) (a)**  $\overline{AB}$ ,  $\overline{Ac}$  are two chords in a circle M. m ( $\angle$ BAC) = 120°. X, Y are the mid points of  $\overline{AB}$ ,  $\overline{AC}$  respectively. Draw  $\overline{XM}$  to cut the circle at D and  $\overline{YM}$  to cut the circle at E. Prove that DE = r. (such r is the radius of the circle)


#### (b) In the opposite figure

 $\overrightarrow{XY}$  ,  $\overrightarrow{XZ}$  are two tangent to the circle from the

Point X, m ( $\angle$ D) = 110°

Prove that:

$$m(\widehat{ZE}) = m(\widehat{ZY})$$



**(5) (a)**  $\overline{AB}$  is a diameter of a circle M,  $\overline{AC}$  is a chord, E is the mid point of  $\overline{AC}$ , the tangent  $\overline{BD}$  is drawn to cut  $\overline{AC}$  at D, then draw  $\overline{EM}$  to cut the circle at X. Prove that:

[1] MEDB is cyclic quadrilateral

[2] 2 m (
$$\angle$$
BAX) = m ( $\angle$ D)


#### (b) In the opposite figure

 $\overline{AB}$  ,  $\overline{AC}$  are two chords in a circle M

D , E are the two mid points of  $\overline{AB}$  ,  $\overline{AC}$ 

respectively and m ( $\angle BAC$ ) =  $55^{\circ}$ 

Find m (∠DME)



# Model (3)

# [1] Complete the following

(1) The line of centres of two touching internally circles passing ............


(2) A circle of radius, A is point on its plan, if MA =  $\frac{3}{4}$  R, then A lies .......

(3) if surface of the circle  $M_1 \cap$  the surface of the circle  $M_2 = \emptyset$ , then the two circle are ......

# (4) In the opposite figure

M (
$$\angle$$
ABC) = 63°, m ( $\angle$ CED) = 67°

, then m  $(\angle C) = \dots$ 




(5) The two opposite angles of be cyclic quad are ..........

#### (6) In the opposite figure

$$m(\widehat{CE}) = 100^{\circ}$$
,  $m(\angle \widehat{BD}) = 30^{\circ}$ 

then m (∠A) = ......



#### [2] Choose the correct from the given ones:

- (1) The ratio between the measure of the inscribed angle and the measure of the central angle subtended by the same arc equals .......
  - (a) 1:2
- (b) 2:1
- (c) 1:1
- (d) 1:3
- (2) The measure of the arc which represents  $\frac{1}{3}$  the measure of the circle equals.......
  - (a) 60°
- (b) 90°
- (c) 120°
- (d) 240°

- (3) If  $R_1$ ,  $R_2$  are the radii of the two circles M,N and MN>  $R_1+R_2$ , them the two circles are .......
  - (a) touching internally
- (b) touching externally

(c) intersecting

- (d) distant
- (4) If the two circles M, N are touching externally , the length of one radius of them is 3cm, MN = 7cm, then the length of the radius of the circle is
  - (a) 3cm
- (b) 4cm
- (c) 7cm
- (d) 10cm
- (5) A chord of length 8cm in a circle of diameter 10cm, then chord at ....... from the cente of the circle .......
  - (a) 3cm

(b) 4cm

(c) 7cm

- (d) 10cm
- (6) In the opposite figure:

- $(a) 20^{\circ}$
- (b) 40°
- $(c) 60^{\circ}$
- (d) 80°

# (3) (a) In the opposite figure:


Two concentric circles of center M,

 $\overrightarrow{EC}$  is a tangent to the greatest circle,

 $\overline{EB}$  cuts the smallest circle at A, B.


D is the mid point of  $\overline{AB}$  and m ( $\angle$ CED) = 40°.

Find with proof m ( $\angle DMC$ ).




#### (b) In the opposite figure


In the circle M, m ( $\angle$ BMC) = 100°, m ( $\angle$ ABD) = 120°. Find with proof m ( $\angle$ DCB).





# (4) (a) In the opposite figure


M and N are two distant and congruent Circles. Draw  $\overrightarrow{AD}$  intersecting circle M at A and B intersects circle N at C and D respectively, if  $\overline{MX} \perp \overline{AB}$ ,  $\overline{NY} \perp \overline{CD}$ , F is the midpoint of  $\overline{MN}$ .



















solution set of the equation  $x^2 - 4x + 3 = 0$  and write the equation of the axis of symmetry

(b)If A ,B are two events in a random experiment, P(A) =0.7 , P(B) = 0.6 and  $P(A \cap B)$  = 0.4 , find the probability of :

- (1) Non occurrence of the event A.
- (2) occurrence of one event without the other.

